

# TWO-DIMENSIONAL MHD NUMERICAL SIMULATIONS OF MAGNETIC RECONNECTION TRIGGERED BY A SUPERNOVA SHOCK IN INTERSTELLAR MEDIUM: GENERATION OF X-RAY GAS IN GALAXY

SYUNITI TANUMA <sup>1,2,3</sup>, TAKAAKI YOKOYAMA, <sup>4</sup>, TAKAHIRO KUDOH <sup>2,5</sup>,  
AND

KAZUNARI SHIBATA <sup>6</sup>

*Draft version February 1, 2008*

## ABSTRACT

We examine the magnetic reconnection triggered by a supernova (or a point explosion) in interstellar medium, by performing two-dimensional resistive magnetohydrodynamic (MHD) numerical simulations with high spatial resolution. We found that the magnetic reconnection starts long after a supernova shock (fast-mode MHD shock) passes a current sheet. The current sheet evolves as follows: (i) Tearing-mode instability is excited by the supernova shock, and the current sheet becomes thin in its nonlinear stage. (ii) The current-sheet thinning is saturated when the current-sheet thickness becomes comparable to that of Sweet-Parker current sheet. After that, Sweet-Parker type reconnection starts, and the current-sheet length increases. (iii) “Secondary tearing-mode instability” occurs in the thin Sweet-Parker current sheet. (iv) As a result, further current-sheet thinning occurs and anomalous resistivity sets in, because gas density decreases in the current sheet. Petschek type reconnection starts and heats interstellar gas. Magnetic energy is released quickly while magnetic islands are moving in the current sheet during Petschek type reconnection. The released magnetic energy is determined by the interstellar magnetic field strength, not energy of initial explosion nor distance to explosion. We suggest that magnetic reconnection is a possible mechanism to generate X-ray gas in Galaxy.

*Subject headings:* magnetic reconnection — Galaxy: structure — ISM: magnetic fields — MHD — shock waves — X-rays: ISM

## 1. INTRODUCTION

Observations of the diffuse component of X-rays (e.g. Snowden 1995) indicate that hot components of temperatures higher than  $10^6$  K occupy a considerable volume of the interstellar space. These hot components may be generated by supernova remnants (SNRs), stellar winds and magnetic heating. X-rays from Galactic ridge (GRXE; Galactic Ridge X-ray Emission; see discussion; Kaneda et al. 1997; Koyama et al. 1986a; Makishima 1994; Yamasaki et al. 1997; Yamauchi et al. 1996; Valinia & Marshall 1998), for example, can not be fully explained by the point sources such as SNRs and stellar winds (Kaneda et al. 1997). We study the magnetic heating in interstellar

space in this paper. The mean strength of interstellar magnetic field is several  $\mu\text{G}$  (Beck et al. 1996; Heiles et al. 1976; Sofue et al. 1986; Vallée 1997). The interstellar magnetic field has random components and generates possibly many current sheets at all scales (e.g. Ohno & Shibata 1993). Parker (1992) pointed out the importance of magnetic reconnection for the heating of Galactic plasma (see also Hanasz & Lesch 1998). Magnetic reconnection is a fundamental intrinsic property of agitated magnetized, turbulent plasma. Whenever the magnetic fields collide with another field with different direction by, for example, the shear motion, magnetic energy is rapidly dissipated. Outside the forming current sheet the mag-

<sup>1</sup> Department of Astronomy, School of Science, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

<sup>2</sup> National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka-shi, Tokyo 181-8588, Japan

<sup>3</sup> Present Address: Solar-Terrestrial Environment Laboratory, Nagoya University, 3-13 Honohara, Toyokawa, Aichi 442-8507, Japan : tanuma@stelab.nagoya-u.ac.jp

<sup>4</sup> Nobeyama Radio Observatory, Minamimaki, Minamisaku, Nagano 384-1305, Japan : yokoyama@nro.nao.ac.jp

<sup>5</sup> kudoh.takahiro@nao.ac.jp

<sup>6</sup> Kwazan Observatory, Faculty of Science, Kyoto University, Ominemachi, Kitakazan, Yamashina-ku, Kyoto 607-8471, Japan : shibata@kwasan.kyoto-u.ac.jp

netic field lines are frozen-in to the interstellar gas, which is provided by the high electrical conductivity. When the oppositely directed field lines collide with each other, the field gradient steepens and the current density increases until strong dissipation sets in (e.g. via anomalous resistivity) to trigger fast reconnection (e.g. Hanasz & Lesch 1998; Ugai 1986). The reconnection heats the interstellar gas by releasing interstellar magnetic energy, and accelerates it by magnetic tension force to Alfvén velocity, forming interstellar jets. The magnetic reconnection plays an important role as observed in solar flares by X-ray satellite *Yohkoh* (Tsuneta 1996; Shibata 1996), and in substorms by satellite such as *Geotail* (e.g. Hoshino et al. 1998). In the solar atmosphere, the magnetic reconnection heats the plasma from a temperature of several  $\times 10^6$  K to several  $\times 10^7$  K (sometimes several  $\times 10^8$  K), and accelerates it to  $10^{2-3}$  km s $^{-1}$  (e.g. Shibata 1996; Yokoyama & Shibata 1995). If an explosion such as a supernova compresses the current sheet, magnetic reconnection occurs and heats interstellar gas.

Historically, the steady reconnection mechanisms were proposed on the basis of analytical studies. In the Sweet(1958)-Parker(1957) type reconnection, the diffusion region is so long as to occupy whole current system. Therefore, the stored magnetic energy is released by Ohmic heating, which can hardly be applied to solar flare phenomena, because the reconnection rate of the Sweet-Parker model is too small ( $\sim R_m^{-1/2}$ ) in the solar corona, where  $R_m$  is magnetic Reynolds number ( $\sim 10^{10-15}$ ). On the other hand, in the Petschek(1964) type reconnection, the diffusion region is localized near an X-point, and standing slow shocks occupy whole current systems. In the configuration, the motor effect associated with slow shocks is much more dominant than Ohmic heating, which can hence be applicable to solar flare phenomena, because the reconnection rate of the Petschek model is  $\sim 0.1 - 0.01$ , independent of  $R_m$  when  $R_m$  is large. This is called “fast reconnection”. Two models have been suggested to clarify how the fast reconnection can be realized. One is so-called “externally driven fast reconnection model”, which predicts that the fast reconnection should be controlled by boundary conditions (Petschek 1964; Sato & Hayashi 1979; Priest & Forbes 1986). The other is so-called “spontaneous fast reconnection model”, which predicts that the fast reconnection mechanism spontaneously develops from inside the system by self-consistent interaction between plasma microscopic processes (current-driven anomalous resistivities) and macroscopic reconnection flows (see also Ugai 1986).

The purpose of this paper is to examine the magnetic reconnection triggered by a supernova by per-

forming two dimensional (2D) numerical magnetohydrodynamic (MHD) simulations including the electric resistivity. Many MHD numerical simulations have been performed for magnetic reconnection in the solar atmosphere (e.g. Yokoyama & Shibata 1995; Odstrčil & Karlický 1997). Some similar attempts have been made in the model of intracluster medium (Valinia et al. 1996; see also Tajima & Shibata 1997), Galactic halo (Birk, Lesch, & Neukirch 1998; Tanuma et al. 1998; Zimmer, Lesch, & Birk 1997), and protostar (Hayashi, Shibata, & Matsumoto 1996; Shibata & Yokoyama 1999). The magnetic reconnection occurs at numerous current sheets in Galactic disk and halo, which may be called “galactic flare” (Sturrock & Stern 1980). Recently, Tanuma et al. (1999) performed numerical simulations and suggested that the magnetic reconnection can generate the Galactic-ridge X-ray emission (see section 4.2). In this paper, we solve the 2D resistive MHD equations numerically to see how the magnetic reconnection occurs when a supernova triggers the magnetic reconnection in Galaxy. In this paper, we analyze the detailed time evolution of the reconnecting current sheet, study dependence of the results on many parameters, and apply the results to Galaxy.

In the next section, we describe the method of the 2D resistive MHD numerical simulations of the magnetic reconnection triggered by a supernova as the simplest model. In section 3, we describe the results of the simulations, and we apply the results to the X-ray plasmas in Galaxy in section 4. In the last section, we summarize this paper.

## 2. METHOD OF NUMERICAL SIMULATION

### 2.1. The Situation of Problem

The magnetic reconnection occurs when the interstellar magnetic field collides with another magnetic field which is not exactly parallel. The magnetic reconnection heats and accelerates interstellar gas by releasing magnetic energy. In the present paper, we treat the magnetic reconnection triggered by a supernova (figure 1). We assume a simple initial condition (uniform temperature everywhere, and uniform magnetic field outside the current-sheet) and large simulation region. We assume the magnetic fields are anti-parallel to each other, and a supernova occurs near the current sheet. We solved the two-dimensional (2D) resistive magnetohydrodynamic (MHD) equations numerically to see how the magnetic reconnection heats the interstellar gas.

### 2.2. MHD Basic Equations

The 2D resistive MHD basic equations are written as follows:

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{v}) = 0, \quad (1)$$

$$\rho \frac{\partial \mathbf{v}}{\partial t} + \rho (\mathbf{v} \cdot \nabla) \mathbf{v} + \nabla p_g = \frac{1}{c} \mathbf{J} \times \mathbf{B} + \rho \mathbf{g}, \quad (2)$$

$$\frac{\partial \mathbf{B}}{\partial t} - \nabla \times (\mathbf{v} \times \mathbf{B}) = -c \nabla \times (\eta \mathbf{J}), \quad (3)$$

$$\frac{\partial e}{\partial t} + \nabla \cdot [(e + p_g) \mathbf{v}] = \eta |\mathbf{J}|^2 + \mathbf{v} \cdot \nabla p_g \quad (4)$$

where  $\rho$ ,  $\mathbf{v}$ ,  $\mathbf{B}$ ,  $\eta$ ,  $e$ ,  $\mathbf{J}$ , and  $\mathbf{g}$  are mass density, velocity, magnetic field, electric resistivity, internal energy, current density ( $= c \nabla \times \mathbf{B} / 4\pi$ ), and gravity ( $= 0$ ), respectively. We use the equation of state for the ideal gas, i.e.,  $p_g = (\gamma - 1)e$  where  $\gamma$  is the specific heat ratio ( $= 5/3$ ).

### 2.3. Normalization

We normalize length, velocity, and time. The units are

$$H \simeq 100 E_{51}^{1/3} \left( \frac{p_{13}}{3} \right)^{-1/3} \text{ pc}, \quad (5)$$

$$C_s = (\gamma R_g T_0)^{1/2} \simeq 20 T_4^{1/2} \text{ km s}^{-1}, \quad (6)$$

$$\tau = \frac{H}{C_s} \simeq 3 \times 10^6 E_{51}^{1/3} T_4^{-1/2} \left( \frac{p_{13}}{3} \right)^{-1/3} \text{ yr}, \quad (7)$$

respectively, where  $E_{51}$ ,  $T_4$ ,  $p_{13}$ ,  $C_s$ , and  $R_g$  are the initial explosion energy ( $E_{\text{ex}}$ ) in unit of  $10^{51}$  erg, temperature in unit of  $10^4$  K, gas pressure in units of  $10^{-13}$  erg cm $^{-3}$ , sound velocity, and gas constant, respectively. We assume that  $H$  is derived from

$$E_{\text{ex}} = \frac{p_{\text{ex}}}{\gamma - 1} \left( \frac{4\pi}{3} H^2 D_{\text{SN}} \right). \quad (8)$$

where  $D_{\text{SN}} (= H)$  is a half-thickness of assumed initial high-pressure region in a direction perpendicular to  $x$ - $y$  plane, and  $p_{\text{ex}} = p_{\text{ratio}} p_g = 500 p_g = 5 \times 10^{-11} p_{13}$  (Model A1).

The units of density, gas pressure, magnetic field strength, current density, and resistivity are  $\rho_0 \sim 1 \times 10^{-25}$  g cm $^{-3}$ ,  $\rho_0 C_s^2 \sim 4 \times 10^{-13}$  erg cm $^{-3}$ ,  $(\rho_0 C_s^2 / 8\pi)^{1/2} \sim 3.2$   $\mu$ G,  $(\rho_0 C_s^2 / 8\pi)^{1/2} / H$ , and  $c^2 H C_s / 4\pi$ , respectively. In this paper, we solve non-dimensional basic equations, and describe the results of numerical simulations by non-dimensional values.

### 2.4. Anomalous Resistivity Model

We assume the anomalous resistivity model as follows:

$$\eta = \begin{cases} \eta_0 & \text{if } v_d \leq v_c \\ \eta_0 + \alpha (v_d / v_c - 1)^2 & \text{if } v_d > v_c \end{cases} \quad (9)$$

(Ugai 1992; Yokoyama & Shibata 1995), where  $v_d (= J/\rho)$ ,  $\rho$ ,  $J$ ,  $v_c$ , and  $\eta_0$  are the normalized relative ion-electron drift velocity, non-dimensional mass density, non-dimensional current density, threshold above which anomalous resistivity sets in, and “background uniform resistivity”. The parameters are  $\eta_0 = 0.015$ ,  $\alpha = 0.1$  and  $v_c = 100$  in this paper for the typical model (Model A1).

### 2.5. Initial Condition of Typical Model (Model A1)

Figure 2 shows the initial condition of our numerical simulation. We assume the temperature is  $T = T_0 = 1$  (uniform) everywhere. The current-sheet thickness is  $2l^{\text{init}} = 2$ . We take Cartesian coordinate  $(x, y)$ . The magnetic field, gas pressure, and density are

$$\mathbf{B} = B_0 \tanh(y) \mathbf{x}, \quad (10)$$

$$p_g = p_0 + p_{\text{mag}}^{\text{init}} \cosh^{-2}(y), \quad (11)$$

$$\rho = \gamma p_g / T = \rho_0 + (\gamma p_{\text{mag}}^{\text{init}} / T_0) \cosh^{-2}(y), \quad (12)$$

respectively where  $B_0 \simeq 8.68$ ,  $\mathbf{x} = (1, 0)$ ,  $p_0 = 1/\gamma = 0.6$ ,  $\rho_0 = 1$ , and  $p_{\text{mag}}^{\text{init}}$  is initial magnetic pressure ( $= B_0^2 / 8\pi = p_0 / \beta$ ) outside the current sheet. The ratio of gas to magnetic pressure is  $\beta = 8\pi p_0 / B_0^2 = 0.2$  ( $|y| > 1$ ). Total pressure ( $P = p_g + |\mathbf{B}|^2 / 8\pi$ ) is uniform everywhere. The sound velocity is  $C_s \equiv (\gamma p_g / \rho)^{1/2} = 1$  (uniform) in the initial condition. The initial Alfvén velocity is  $v_A^{\text{init}} = B_0 / (4\pi \rho_0)^{1/2} \simeq 2.45$  ( $|y| > 1$ ).

We assume the symmetric boundaries for the top ( $y \simeq 81.7$ ) and bottom ( $y \simeq -54.8$ ) surfaces, and the periodic boundaries for the left ( $x \simeq -120.9$ ) and right ( $x \simeq 120.9$ ) surfaces. The simulation region size is  $L_x \sim 241.8$  times  $L_y \sim 136.5$ . The number of grid points is  $(N_x \times N_y) = (465 \times 602)$ . The grid sizes are non-uniform ( $\Delta x \geq 0.20$ ,  $\Delta y \geq 0.025$ ). The number of grid points in the current sheet is 80. We put a high-pressure region instead of a point explosion (supernova) at  $(x, y) = (0, y_{\text{ex}}) = (0, 7)$ , whose radius and gas pressure are  $r_0 = 1$  and  $p_{\text{ex}} = p_{\text{ratio}} p_g = 500 p_0$ , respectively.

We use the 2-steps modified Lax-Wendroff method. We neglect gravitational force, effects of rotation such as shearing motion and Coriolis force, cosmic rays, viscosity, radiative cooling, and heat conduction in our simulation models. The cooling times due to heat conduction and radiation are

$$\begin{aligned} \tau_{\text{cond}} &\sim \frac{nkT\lambda^2}{\kappa_0 T^{7/2}} \\ &\sim 10^{13} \left( \frac{n}{0.1 \text{ cm}^{-3}} \right) \end{aligned}$$

$$\left(\frac{\lambda}{100 \text{ pc}}\right)^2 \left(\frac{T}{10^4 \text{ K}}\right)^{-5/2} \text{ yr}, \quad (13)$$

$$\tau_{\text{rad}} \sim \frac{nkT}{n^2\Lambda(T)}$$

$$\sim 10^3 \left(\frac{T}{10^4 \text{ K}}\right) \left(\frac{n}{0.1 \text{ cm}^{-3}}\right)^{-1} \left[\frac{\Lambda(10^4 \text{ K})}{10^{-21} \text{ erg cm}^3 \text{ s}^{-1}}\right]^{-1} \text{ yr}, \quad (14)$$

respectively, for cool, dense gas, where  $\Lambda$  is the cooling function (Spitzer 1962), and  $\kappa_0$  is constant ( $= 10^{-6} \text{ erg s}^{-1} \text{ cm}^{-1} \text{ K}^{-1}$ ). The radiative cooling time becomes  $\tau_{\text{rad}} \sim 10^6 (n/0.1 \text{ cm}^{-3})^{-1} \text{ yr}$  for the hot gas of  $T \sim 10^7 \text{ K}$ , where  $\Lambda(10^7 \text{ K})$  is  $\sim 10^{-23} \text{ erg cm}^3 \text{ s}^{-1}$ . Furthermore, in the X-ray gas near Galactic plane (Galactic Ridge X-ray Emission=GRXE;  $T \sim 10^8 \text{ K}$  and  $n \sim 3 \times 10^{-3} \text{ cm}^{-3}$ ; see Discussion), the radiative cooling time is  $\sim 10^{10.5} \text{ yr}$ , which is much longer than the typical time scale ( $3 \times 10^6 \text{ yr}$ ; see next section) of the physical process examined in this paper, so that the radiative cooling can be neglected. On the other hand, the conduction cooling time for the X-ray gas is  $\sim 30 \text{ yr}$  [which is  $\sim 10^5 (\lambda_{\text{eff}}/1 \text{ kpc})(T/10^8 \text{ K})^{-1/2} \text{ yr}$  in fact because GRXE gas is collisionless]  $\ll 3 \times 10^6 \text{ yr}$ , so that the conduction cooling can not be neglected, where  $\lambda_{\text{eff}}$  is the effective length of helical magnetic field. Nevertheless, we neglect the effect of heat conduction for simplicity in this paper, since the basic properties of magnetic reconnection such as reconnection rate nor energy-release rate are not much affected by the heat conduction (Yokoyama & Shibata 1997). We assume so long current sheet that the effect of evaporation is negligible, though the evaporation occurs if the interstellar cloud exists near the reconnection region and the reconnected field lines penetrate the cloud. In this paper, we assume the uniform strong field and long current sheet under pressure equilibrium for simplicity, instead of a small turbulence or small current sheets under non-equilibrium.

### 3. RESULTS

#### 3.1. Typical Model (Model A1)

Figure 3 shows the time evolution of the two-dimensional distribution of the current density, density, and gas pressure, with magnetic field lines, in an early phase of the interaction between the supernova and current sheet. A supernova occurs near the current sheet. A blast shock passes and perturbs the current sheet (figure 3). Figures 4-6 show the time evolution of spatial distributions of the temperature (figure 4), gas pressure (figure 5), and current density (figure 6), with velocity vectors and magnetic field lines. Figure 7 shows the schematic illustration of simula-

tion results. The current sheet evolves and magnetic reconnection occurs sequentially as the following 5 phases. (i) Shock wave passage ( $t < 2$ ; figure 7[a]) and the current sheet thinning by the tearing instability in nonlinear phase ( $t \sim 100 - 200$ ; figure 7[b]). (ii) Sweet-Parker type reconnection ( $t \sim 200 - 270$ ; figure 7[c]). (iii) “Secondary tearing instability” in the Sweet-Parker current sheet ( $t \sim 250 - 270$ ; figure 7[d]). (iv) Generation of magnetic islands ( $t \sim 270$ ; figure 7[e]) and Petschek type (fast) reconnection with two standing slow shock regions ( $t \sim 270 - 400$ ; figure 7[f]). (v) End of Petschek type reconnection ( $t \sim 400$ ).

The current sheet is unstable to the tearing mode instability (Furth, Killeen, & Rosenbluth 1963). The magnetic dissipation time, Alfvén time, and time scale of the tearing instability in the initial state are

$$\tau_{\text{dis}}^{\text{init}} = \frac{l^{\text{init}2}}{\eta_0} \sim \frac{1^2}{0.015} \sim 67, \quad (15)$$

$$\tau_A^{\text{init}} = \frac{l^{\text{init}}}{v_A^{\text{init}}} \sim \frac{1}{2.5} \sim 0.4, \quad (16)$$

$$\tau_t^{\text{init}} = (\tau_{\text{dis}}^{\text{init}} \tau_A^{\text{init}})^{1/2} \sim 5.2, \quad (17)$$

respectively, where  $l^{\text{init}} (= 1)$  is the half-thickness of initial current sheet. The magnetic Reynolds (Lundquist) number is

$$R_{\text{m,y}}^{\text{init}} \equiv \frac{v_A^{\text{init}} l^{\text{init}}}{\eta_0} \sim 167. \quad (18)$$

**Phase I: Shock Wave Passage ( $t < 2$ ) and Current-Sheet Thinning by Tearing Instability ( $t \sim 100 - 200$ )** The supernova blast shock (a fast-mode magnetohydrodynamic shock) crosses the current sheet in an early phase ( $t < 2.5$ ; figure [3]). The magnetic reconnection does not occur immediately after the shock wave crosses the current sheet, because the crossing time is too short to drive the magnetic reconnection.

Figure 8 shows the time variation of half-thickness of the current sheet, defined by the minimum half-width of the half-maximum of gas pressure calculated parallel with  $y$ -axis in  $|x| < 25$ . Figure 9 shows its schematic illustration. The magnetic field is dissipated ( $t \sim 0 - 100$ ), and the current-sheet thickness increases slowly. The current-sheet thinning occurs by the tearing instability ( $t \sim 100 - 200$ ). Figure 10 shows the time variation of magnetic flux, defined by  $\psi(t) = \int_0^t \max_{|x| < 10} [(B_x v_y)[\text{at}(x, 2)] + (B_x v_y)[\text{at}(x, -3)]] dt$ . The process in this phase corresponds to the nonlinear stage of the tearing-mode instability (Furth, Killeen, & Rosenbluth 1963; Magara & Shibata 1999; Steinolfson and & Hoven 1983,

1984). The growth rate is consistent with the results of Magara & Shibata (1999). The dashed line in figure 10 shows  $\log \psi(t) \sim 0.00257t$ . It is  $1/2$  times that of the linear phase [ $\log \psi(t) = \log_{10} e \times 0.43(1/\tau_t^{\text{init}})(1/4\pi)t \sim 0.0042t$ ; Magara & Shibata 1999]. It explains the result of our numerical simulation well.

The half-thickness of the current sheet decreases to  $l \sim 0.35$  at  $t \sim 200$  when the current-sheet thinning stops. The current-sheet length is comparable to the most unstable wavelength of the tearing instability occurring in the initial current sheet, i.e.,

$$\lambda_t \sim 5.6 R_{m,y}^{\text{init} 1/4} l^{\text{init}} \quad (19)$$

(Magara & Shibata 1999). It is  $\sim 20$  in this model. The current-sheet thickness in this phase is explained by

$$l_t \sim \left(\frac{\lambda_t}{2}\right) R_{m,t}^{-1/2}, \quad (20)$$

i.e., this current sheet corresponds to Sweet-Parker current sheet (Magara & Shibata 1999). Here, the magnetic Reynolds number is

$$R_{m,t} = \frac{(\lambda_t/2)v_A^{\text{init}}}{\eta_0}. \quad (21)$$

The equations (20) and (21) give us  $l_t \sim 0.23$  and  $R_{m,t} \sim 1467$ , respectively. These values explain our results in this phase well. In summary, we can say that the current-sheet thinning stops when the current sheet evolves to Sweet-Parker current sheet. The gas flows into the diffusion region at the velocity of  $v_{\text{in}} \sim 0.01$ . The duration of this phase is  $\Delta t_t \sim 100$ , which is approximately equal to the time for initial current sheet to become the Sweet-Parker sheet,  $l^{\text{init}}/v_{\text{in}} \sim 1/0.01 \sim 100 \sim 20\tau_t^{\text{init}}$ .

**Phase II: Sweet-Parker Type Reconnection ( $t \sim 200 - 270$ )** Sweet(1958)-Parker(1957) type reconnection starts at  $t \sim 200$  in an asymmetric situation. Alfvén velocity is higher in the region ( $y > 0$ ) because of the low density gas generated by the supernova. The gas flows along current sheet, and is accelerated to  $v \sim 2.5$ , which is equal to the Alfvén velocity near  $(x, y) \sim (0, -2)$ . Figure 11 shows (a) the time variation of velocity of the gas flowing into the reconnection region. The inflow velocity is defined by  $v_{\text{in}} = \max_{|x| < 10} [v_y[\text{at}(x, -3)] - v_y[\text{at}(x, 1)]] / 2$ . Figure 11(b) shows Alfvén velocities attained at  $(x_0, 2)$  (the solid line) and at  $(x_0, -2)$  (the dashed line) where  $x_0$  is the point where maximum inflow velocity attained. Figure 11(c) shows the reconnection rate defined by  $v_{\text{in}}/v_{A,s}$ , where  $v_{A,s}$  is the smaller one between  $v_A[\text{at}(x_0, 1)]$  and  $v_A[\text{at}(x_0, -3)]$ . The

current-sheet length increases to  $\lambda \sim 80$  (figure 8), because the gas in the initial current sheet flows out along the current sheet. The inflow velocity becomes  $v_{\text{in}} \sim 0.025$ . The ratio of inflow to outflow velocity ( $\sim 0.025/2.5 \sim 0.01$ ) is explained by

$$\frac{v_{\text{in}}}{v_A} \sim \frac{l_{\text{SP}}}{\lambda_{\text{SP}}/2} \sim R_{m,\text{SP}}^{-1/2} \quad (22)$$

where  $R_{m,\text{SP}}$  is the local magnetic Reynolds number defined by

$$R_{m,\text{SP}} = \frac{(\lambda_{\text{SP}}/2)v_A}{\eta_0}, \quad (23)$$

$\lambda_{\text{SP}}$  and  $l_{\text{SP}}$  are the length and half-thickness of the Sweet-Parker current sheet (Parker 1957), respectively, at the time when the secondary tearing instability starts. These equations give us  $v_{\text{in}}/v_A \sim 0.01$  with  $R_{m,\text{SP}} \sim 10^4$ ,  $\lambda_{\text{SP}} \sim 80$ , and  $l_{\text{SP}} \sim 0.3$ , respectively, in this model. The condition for the setup of the secondary tearing instability is determined by the aspect ratio of Sweet-Parker current sheet, so that the secondary tearing instability occurs when aspect ratio [length/thickness =  $(\lambda/2)/l$ ] becomes larger than 100 (Biskamp 1993). The equation (22) explains the result of our simulation in this phase ( $t \sim 220 - 250$ ) well. During the Sweet-Parker type reconnection, the current-sheet thickness increases in proportion to  $\lambda^{1/2}$ , which is derived from the equation (22).

At  $t \sim 200 - 220$ , the inflow velocity ( $v_{\text{in}}$ ) increases to  $\sim 0.15$  (see the arrow in figure 11), which is much larger than Sweet-Parker value  $\sim R_{\text{SP}}^{-1/2} v_A \sim 0.025$ . This transient inflow velocity is determined by  $v_{\text{in}} \sim 2l_{210}v_{\text{out}}/\lambda_t \sim 0.15$ , which is derived from the mass conservation equation  $\lambda_t v_{\text{in}} \sim 2l_{210}v_{\text{out}}$ , assuming the density is constant and  $v_{\text{out}} (\sim 0.6)$  is the velocity of the outflowing gas along the current sheet at this time ( $t \sim 210$ ), where  $l_{210} \sim 2.5$  is the current-sheet half-thickness in  $x \sim 20$  (at the edge of Sweet-Parker current sheet) at  $t \sim 210$  (Cf. Fu & Lee 1986; Shibata 1997).

Figure 12 shows the time variation of various energies in the reconnection region ( $|x| < 25$  and  $|y| < 3$ ). The magnetic energy (the solid line), thermal energy (the dotted line), and kinetic energy (the dashed line) are defined by  $E_{\text{mag}} = \int_{-25}^{25} \int_{-3}^3 (B^2/8\pi) dx dy$ ,  $E_{\text{th}} = \int_{-25}^{25} \int_{-3}^3 [p_g/(\gamma - 1)] dx dy$ , and  $E_{\text{kin}} = \int_{-25}^{25} \int_{-3}^3 (\rho v^2/2) dx dy$ , respectively. They are corrected by the energy flux at the boundaries ( $|x| = 25$  and  $|y| = 3$ ). Figure 13 shows the schematic illustration of the time variation of the magnetic energy-release rate. The magnetic energy is converted to the thermal energy slowly by the tearing instability and Sweet-Parker type reconnection. At  $t \sim 200 - 220$ , the magnetic energy-release rate increases (see the ar-

rows in figures 12 and 13) by the nonlinear growth of tearing instability.

**Phase III: Secondary Tearing Instability ( $t \sim 250 - 270$ )** The Alfvén time and time scale of the tearing instability of the Sweet-Parker current sheet become,

$$\tau_{A,SP} = \frac{l_{SP}}{v_A} \sim 0.12, \quad (24)$$

$$\tau_{t,SP} = \left(\frac{l_{SP}^3}{\eta_0 v_A}\right)^{1/2} \sim 0.85, \quad (25)$$

respectively. The tearing instability (we call “secondary tearing instability” in this paper) occurs in Sweet-Parker current sheet at  $t \sim 250$ . The current-sheet thickness decreases again. Some magnetic islands are generated with intervals of  $\sim 5 - 10$  in  $x$ , which are determined by the wavelength of secondary tearing instability,

$$\lambda_2 \sim 4.9 R_{m,y}^{1/4} l_{SP}, \quad (26)$$

where  $R_{m,y}$  is the local Reynolds number, defined by

$$R_{m,y} = \frac{v_A l_{SP}}{\eta_0}, \quad (27)$$

and  $v_A$  is the local Alfvén velocity. The equation (26) gives us  $\lambda_2 \sim 4$  with  $R_{m,y} \sim 50$ , which explains the results of our simulation in this phase well. Some small islands collide with one another, and are coalesced to become big islands. The secondary tearing instability goes on for  $\Delta t_{2t} \sim 20$ . It is equal to  $2.5\tau_{t,SP} \sim 20$ .

**Phase IV: Petschek Type Reconnection ( $t \sim 270 - 400$ )** Figure 14 shows the time variations of (a) maximum drift velocity ( $v_d \equiv J/\rho$ ) in the current sheet ( $|x| < 5$  and  $-2.6 < z < 1.0$ ), (b) density where maximum  $v_d$  is attained, (c) current density ( $J$ ) where maximum  $v_d$  is attained, and (d) maximum magnetic reconnection rate defined by  $\eta J$ . The density decreases from 0.5 to 0.1 at the center of the current sheet (figure 14) because the gas flows away along the current sheet from the dissipation region. The maximum drift velocity exceeds the threshold ( $v_c = 100$ ) in the reconnection region immediately after the big magnetic island moves away. The anomalous resistivity sets in at  $t \sim 270$ . The half-thickness of the current sheet decreases to  $l \sim 0.15$  (figure 8). The diffusion region is much localized near the X-point. Petschek (1964) type reconnection begins with various violent phenomena. The gas flows into reconnection region at  $v_{in} \sim 0.1 - 0.2$  (figure 11). The magnetic reconnection rate (figure 11) increases to

$$\epsilon = \frac{v_{in}}{v_A} \sim 0.05 - 0.10, \quad (28)$$

while it is  $\sim 0.02$  during Sweet-Parker type reconnection. In this simulation, the energy-release rate during Petschek type reconnection is only a few times higher than that during Sweet-Parker type reconnection. The latter, however, is much small in the actual Galaxy because it is determined by the magnetic Reynolds number as  $\epsilon = v_{in}/v_a = R_{in}^{-1/2}$ . The gas in the diffusion region are accelerated along the current sheet by magnetic tension force to  $v \sim 2.5 - 3$ , which is comparable to the Alfvén velocity in the region where the maximum velocity of reconnection jet is attained.

The magnetic energy is converted to the thermal energy slowly during the tearing instability and Sweet-Parker type reconnection, and quickly during Petschek type reconnection, particularly while magnetic islands are moving along the current sheet from the reconnection region (figure 12). The energy conversion is suppressed when the magnetic islands are staying in the diffusion region ( $t \sim 310$  and  $360$ ). Total released magnetic energy is  $\sim 5500$  in the reconnection region ( $|x| < 25$  and  $|y| < 3$ ; figure 12), which is much higher than the initial thermal energy of the supernova ( $E_{ex} \sim 1440$ ). The maximum energy release rate and maximum heating rate are  $-dE_{mag}/dt \sim dE_{th}/dt \sim 70 - 80$  at  $t \sim 340$ . They are determined by the Poynting flux entering the reconnection region,

$$-\frac{dE_{mag}}{dt} \sim 2 \frac{B^2}{4\pi} S v_{in}, \quad (29)$$

where  $S$  is the reconnection region size. The above equation gives us  $\sim 80$ , which explains the results of the simulation (The reconnection region size is derived as  $S \sim 37$  if we use the values of  $B \sim 8$  and  $v_{in} \sim 0.2$ ).

The current density increases in the current sheet (at  $t = 210$  and  $250$ ; figure 6) and behind the moving magnetic islands (at  $t = 275$  and  $300$ ; figure 6). The slow shocks are formed behind the moving islands. Two standing slow shocks are generated in the current sheet (at  $t = 325$  and  $350$ ; figure 6). Figure 15 shows the profile of the various physical quantities in  $x = -12$  at  $t = 325$ . The gas pressure, temperature, density, and velocity are high in the current sheet. The magnetic field is weak in the current sheet. The current density is large at the slow shock regions so that the profile of current density has double peaks. It is noted that this Petschek-type reconnection is very time-dependent; Some magnetic islands are repeatedly generated and ejected in the current sheet, and magnetic energy-release rate is controlled by the magnetic islands.

**Phase V: End of Fast (Petschek Type) Reconnection ( $t \sim 400$ )** The current-sheet length is limited by the high gas pressure near the right and left symmetric boundaries ( $x = -120.8$  and  $x = 120.8$ ). Figure 16 shows the distribution of gas pressure in whole simulation region. The current sheet, whose half-thickness was  $l^{\text{init}} = 1$  at  $t = 0$ , expands near the left and right boundaries. The magnetic loops are accumulated at the left and right boundaries so that current-sheet thickness increases from  $\sim 2$  at  $t = 160$  (figure 16[a]) to  $\sim 20$  at  $t = 300$  (figure 16[b]), and eventually to  $\sim 50$  at  $t = 450$  (figure 16[c]). The gas pressure increases in the reconnected loop. The current-sheet length decreases to  $\lambda \sim 50$  at  $t \sim 450$  (figure 16[c]). The ejection from the diffusion region is suppressed by the high gas pressure in the loop near the boundaries, so that the global field configuration tends to be the potential field produced by the localized currents near the boundaries. On the other hand, the magnetic tension force suppresses the inflow to reconnection region because the potential magnetic field outside the current sheet is bent. Consequently, the magnetic reconnection rate decreases suddenly at  $t \sim 380$ . Duration of the Petschek type reconnection is  $\Delta t_P \sim 170$ .

### 3.2. Parameter Survey

#### 3.2.1. Dependence of Results on Magnetic Field Strength ( $\beta$ )

We examine the dependence of results on the magnetic field strength [ $B_0 = (8\pi p_0/\beta)^{1/2}$ ], i.e.,  $\beta$  (Model A2 and B1-5). The time variations of magnetic energy-release rates are shown in figure 17 by the dotted line (Model B1;  $\beta = 0.1$ ), solid line (Model A2;  $\beta = 0.2$ ), and dashed line (Model B2;  $\beta = 0.3$ ). The magnetic energy-release rate decreases with  $\beta$  as  $\sim 250$  ( $\beta = 0.1$ ),  $\sim 70$  ( $\beta = 0.2$ ), and  $\sim 20$  ( $\beta = 0.3$ ), respectively.

Figure 18 (a) shows the  $\beta$ -dependence of maximum magnetic energy release rates during Petschek type reconnection (the solid line) and during Sweet-Parker type reconnection (the dotted-line) in the reconnection region ( $|x| < 25$  and  $|y| < 3$ ). The energy-release rate decreases with  $\beta$ . The rate during Petschek type reconnection is determined by the Poynting flux entering into reconnection region,

$$-\frac{dE_{\text{mag}}}{dt}|_P \sim 2\frac{B^2}{4\pi}Sv_{\text{in}} \propto \beta^{-3/2}, \quad (30)$$

where  $S$  is the reconnection region size,  $v_{\text{in}} = \epsilon v_A = \epsilon B/(4\pi\rho)^{1/2}$  is the inflow velocity to the reconnection region, and  $\epsilon$  is the reconnection rate. This theoretical  $\beta$ -dependence (30) is also shown in the figure 18(a) by the dashed line [ $\sim 2(B_0^2/4\pi)S\epsilon v_A^{\text{init}} \sim 6.8\beta^{-3/2}$

where  $S$  is derived to be  $\sim 26$  if we use the results of the typical model]. It explains the numerical results well. On the other hand, the magnetic energy-release rate during Sweet-Parker reconnection is given by

$$-\frac{dE_{\text{mag}}}{dt}|_{\text{SP}} \sim 2\frac{B^2}{4\pi}Sv_{\text{in}} \propto \beta^{-5/4}, \quad (31)$$

where  $S$  is the reconnection region size,  $v_{\text{in}} = R_{\text{m,SP}}^{-1/2}v_A \propto v_A^{1/2}$  is the inflow velocity to the reconnection region. This theoretical  $\beta$ -dependence (31) is also shown in the figure 18(a) by the dashed-dotted line [ $\sim 2(B_0^2/4\pi)SR_{\text{m,SP}}^{-1/2}v_A^{\text{init}} \sim 2.5\beta^{-5/4}$  where  $S$  is derived to be  $\sim 67$  if we use the results of the typical model]. It explains the numerical results well.

Figure 18(b) presents the times when maximum magnetic energy release rates are attained during Petschek (the solid line) and Sweet-Parker (the dashed line) type reconnections. The time when Sweet-Parker type reconnection starts is determined by the time scale of dissipation in an early phase ( $\sim \Delta t_t$ ) plus the time scale of tearing instability ( $\sim \Delta t_t$ ). It is almost equal to  $\sim 2\Delta t_t$ . The time scale of tearing instability ( $\tau_t \propto v_A^{-1/2} \propto \beta^{1/4}$ ) is presented in figure 18(b) by the dashed-dotted line ( $\sim 230\beta^{1/4}$ ). The time when Petschek type reconnection starts is determined by the time when Sweet-Parker type reconnection starts ( $\sim 2\Delta t_t$ ) plus the time for the Sweet-Parker current sheet to get long until it becomes unstable to the secondary tearing instability,  $\lambda/2 \geq 100l_{\text{SP}} \sim 30$  (Biskamp 1993), which is equal to  $\sim \lambda_{\text{SP}}/v_A$ . The Alfvén time ( $\tau_A \propto v_A^{-1} \propto \beta^{1/2}$ ) is presented in figure 18(b) by the dashed line ( $\sim 580\beta^{1/2}$ ).

Figures 18(c) and (d) show the maximum temperature of heated gas and maximum velocity of the reconnection jet ( $|y| \leq 2$ ). The maximum temperature and the maximum velocity decreases with  $\beta$ . The interstellar gas is heated, by passing the reconnection region, to

$$T_{\text{max}} \sim \left(1 + \frac{1}{\beta}\right) \left(\frac{n_{\text{in}}}{n_{\text{out}}}\right) T_{\text{in}} \propto \left(1 + \frac{1}{\beta}\right), \quad (32)$$

which is derived from the equation  $n_{\text{out}}kT_{\text{max}} \sim n_{\text{in}}kT_{\text{in}} + B^2/8\pi$ . Here, we assume  $(n_{\text{in}}/n_{\text{out}})T_{\text{in}}$  is independent of  $\beta$ . The  $\beta$ -dependence is plotted in figure 18(c) by the dashed line [ $\sim 7(1 + 1/\beta)$ ]. In the simulation, a part of hot gas of the supernova flows into the reconnection region, and re-heated by the reconnection. The gas is accelerated to Alfvén velocity,

$$v_{\text{jet}} \sim v_A = \frac{B}{(4\pi\rho)^{1/2}} \propto \beta^{-1/2}, \quad (33)$$

by the magnetic tension force along the current sheet, which is shown in the figure 18(d) by the dashed line

( $\sim 1.5\beta^{-1/2}$  which is 1.5 times higher than  $v_A^{\text{init}}$  because  $v_A$  increases with time).

### 3.2.2. Dependence of Results on Distance Between Supernova and Current Sheet

We examine the dependence of results on the distance ( $y_{\text{ex}}$ ) between the supernova and current sheet (Models A2 and C1-3). Figures 19(a) and 19(b) illustrate the maximum magnetic energy release rate and the time when it is attained, respectively. The magnetic energy-release rate decreases with  $y_{\text{ex}}$ . The time scale increases with  $y_{\text{ex}}$ . It is because the initial perturbation decreases with  $y_{\text{ex}}$ , the time when magnetic reconnection starts increases and more magnetic field dissipates before the reconnection starts. The maximum temperature (figure 19[c]) decreases with  $y_{\text{ex}}$ . The maximum velocity of reconnection jet (figure 19[d]) does not depend on the distance because it is determined by the Alfvén velocity.

### 3.2.3. Dependence of Results on Initial Energy of Explosion

We put the point explosion with the gas pressure of  $E_{\text{ex}} \propto p_{\text{ex}} = p_{\text{ratio}}p_0$  as the initial condition. We examine the dependence of results on  $p_{\text{ratio}} \propto E_{\text{ex}}$  (Models A2 and D1-5; figure 20). The normalizations given by the equations (5)-(7) change if  $p_{\text{ratio}}$  of the equation (8) changes. We must rewrite the units of length and time as  $H \propto p_{\text{ratio}}^{-1/3}$  and  $\tau \propto p_{\text{ratio}}^{-1/3}$ , respectively. Figure 20(a) presents the maximum magnetic energy release rate. It does not depend on the energy of initial explosion. Figure 20(b) shows the time when maximum magnetic energy release rate is attained. It decreases with  $p_{\text{ratio}}$  because the initial perturbation decreases. Figure 20(c) shows the maximum temperature of gas heated by the reconnection. The temperature increases with  $p_{\text{ratio}}$  because the hot gas of the initial supernova is re-heated by the magnetic reconnection. Figure 20(d) shows the velocity of reconnection jet. It does not depend on  $p_{\text{ratio}}$  because it is determined by the Alfvén velocity.

We examine the model in which two explosions appear symmetrically in both sides [at (0, 7) and (0, -7)] of the current sheet ( $p_{\text{ratio}} = 500$ ; Model D5). The results are plotted in figure 18 by the cross ( $\times$ ). The time scale is much shorter because the initial current sheet is compressed from both sides. Other results are similar to those of typical model.

### 3.2.4. Dependence of Results on Resistivity Model

We examine the dependence of the results on the background uniform resistivity ( $\eta_0$ ) and the resistivity models (either anomalous or uniform for whole evolutions). Figure 21 shows the time variations of

the half-thickness of the current sheet, for Model A2 ( $\eta_0 = 0.015$ ; thick solid line), Model F1 ( $\eta_0 = 0.03$ ; thin dotted line), Model F2 ( $\eta_0 = 0.0075$ ; thin dashed-dotted line), Model F3 ( $\eta_0 = 0.00375$ ; thin dashed-dotted-dotted line), Model F4 ( $\eta_0 = 0.001$ ; thin dashed line), Model F5 ( $\eta_0 = 0.0001$ ; thin solid line), and Model F6 ( $\eta = 0$ ; thick dashed line). In the current sheet of Model A2 ( $\eta_0 = 0.015$ ), the tearing instability ( $t \sim 60 - 150$ ) is followed by Sweet-Parker type reconnection ( $t \sim 150 - 200$ ), secondary tearing instability ( $t \sim 200 - 240$ ), and Petschek type reconnection ( $t \sim 240$ ). In zero-resistivity model (Model F6), the current sheet does not dissipate at first. It, however, starts to become thin at  $t \sim 220$ , and eventually leads to Petschek type reconnection by numerical resistivity (resistivity-like effect by numerical noise). The result of Model F5 ( $\eta_0 = 0.0001$ ) is similar to that of Model F6 ( $\eta = 0$ ), not that of F4 ( $\eta_0 = 0.001$ ). The numerical resistivity in these simulations is estimated as  $\eta_{\text{num}} = 0.001 - 0.0001$ . In Model F1 ( $\eta = 0.03$ ), the current sheet dissipates ( $t < 100$ ), and the magnetic reconnection does not occur ( $t < 800$ ). The maximum temperature of the gas heated by the reconnection and the velocity of the reconnection jet do not depend on  $\eta_0$ .

The result of uniform resistivity model (Model E) is also shown in Figure 18 by triangle ( $\triangle$ ). Petschek type reconnection does not occur in the uniform resistivity model. The magnetic energy-release rate and the time scale are approximately equal to the results of anomalous resistivity model during the Sweet-Parker type reconnection.

### 3.2.5. Dependence of Results of Simulation Region Size and Grid Size

We examine the dependence of results on the simulation region size ( $L_y$ ) in  $y$  axis (Models A2 and G1-2). The magnetic energy release rate increases with the simulation region size because the stored initial total magnetic energy increases. The other results do not depend on the simulation region size.

We examine the effects of grid size ( $\Delta x$ ) (Models A1-2). Until the secondary tearing instability starts, the time evolution of current sheet depends on the parameters such as  $\beta$  and  $\eta$  well so that the time scales of phenomena are predictable. At this stage (the onset of the secondary tearing instability), however, it becomes difficult to predict where small islands appear and when and how they move and collide with one another, since the current-sheet thickness decreases to a few  $\times \Delta y$  after the secondary tearing instability starts. At this stage, the current sheet is too thin to be free from numerical resistivity in our problem. The maximum energy release rate, maximum temperature of the heated gas and velocity of



the reconnection jet do not depend on  $\Delta x$ , although the time when maximum magnetic energy release rate is attained depends on  $\Delta x$  (figure 22).

#### 4. DISCUSSION

Odstrčil & Karlický (1997) performed an interesting 2D numerical simulation of the magnetic reconnection triggered by a point explosion near the current sheet, with parameters different from ours, in order to construct a model of the solar flare triggered by a distant flare. They assumed resistivity as  $\eta \propto |J - J_c|$ , where  $J_c$  is the critical current density above which the anomalous resistivity sets in. Zimmer et al. (1997) tried to explain that the gas is heated in Galactic halo by magnetic reconnection triggered by the collision of the high-velocity clouds (HVC;  $M \sim 10^{7-8} M_\odot$ ;  $V \sim 100 \text{ km s}^{-1}$ ; Blitz 1999), by performing the 2D MHD numerical simulation. Birk et al. (1998) performed 2D numerical simulation of the magnetic reconnection including the ionization and recombination. Above three groups followed the current-sheet thinning and the generation of the magnetic island. They, however, did not reveal the time evolution of the secondary tearing instability, because they did not use enough small grid size. In this paper, we use 80 grid points, so that we revealed how the secondary tearing instability leads to Petschek type magnetic reconnection.

We revealed how the fast reconnection starts at the current sheet via the secondary tearing instability by assuming the anomalous resistivity model (equation [9]) in this paper. Main result of this paper is revealing how the current sheet leads to fast reconnection via the secondary tearing instability. This result does not depend on the form of resistivity before the anomalous resistivity sets in (Yokoyama & Shibata 1995; Ugai 1992). The simulation results we apply to the actual Galaxy, such as temperature of heated gas, do not depend on the resistivity model (Yokoyama & Shibata 1995; Ugai 1992).

We assume that the initial current-sheet has enhanced gas pressure [ $p_g = p_0 + (p_0/\beta) \cosh^{-2}(y)$ ;  $B_x = B_0 \tanh(y)$ ;  $B_y = 0$ ] in the paper. The tearing instability grows similarly in both this model (enhanced gas pressure model) and force-free model [ $p_g = p_0$ ;  $B_x = B_0 \cos 0.5\pi(y + 1)$ ;  $B_z = B_0 \sin 0.5\pi(y + 1)$ ;  $B_y = 0$ ; Magara & Shibata 1999]. The basic physics of magnetic reconnection do not change, though the nonlinear phase starts earlier in the enhanced gas pressure models (Magara & Shibata 1999). The physical processes of our results are hardly influenced by the boundary conditions because the simulation region is large enough. We do not assume the initially enhanced resistivity in the current sheet to see how the spontaneous reconnection oc-

curs after the secondary tearing instability. We use so small grid size that we can suppress the numerical resistivity in our numerical simulations.

The results do not depend on  $p_{\text{ratio}}$  very much. The magnetic reconnection occurs with any  $p_{\text{ratio}}$ . It means that the magnetic reconnection occurs not only by a supernova, but also by the small perturbation. The magnetic reconnection can be triggered by various mechanisms such as a supernova (or a point explosion), superbubbles (sum of supernovae; Tomisaka 1998; Tenorio-Tangle & Bodenheimer 1988), stellar winds, collision of interstellar clouds, shock waves, Parker(1966) instability (undular mode of magnetic buoyancy instability), magnetorotational instability (Balbus & Hawley 1991), collision of galaxies, cosmic rays etc. in numerous current sheets.

The random field stores the magnetic energy at  $\sim$  several 10 pc (Rand & Kulkarni 1989). Total magnetic energy is  $E_{\text{mag}} \sim [\langle B \rangle_{\text{obs}}^2 / 8\pi] V_G \sim 10^{54.4} \text{ erg}$ , where  $\langle B \rangle_{\text{obs}} \sim 3 \mu\text{G}$  is assumed as the mean observed field strength, and  $V_G \sim 2 \times 10^2 \text{ kpc}^3$  is the volume of Galaxy. The thermal energies of GRXE (Galactic Ridge X-ray Emission) and coronal halo are  $E_{\text{GRXE}} \sim 10^{55} \text{ erg}$  (e.g. Kaneda et al. 1997) and  $E_{\text{halo}} \sim 10^{57.2} \text{ erg}$  (Habe & Ikeuchi 1980; Li & Ikeuchi 1992; Ikeuchi 1988), respectively. Total rotational energy of Galaxy ( $E_{\text{rot}} \sim 10^{58.9} \text{ erg}$ ) and total kinetic energy of the interstellar gas ( $E_{\text{ISM}} \sim 10^{58.2} \text{ erg}$ ) are higher than the thermal and magnetic energies (Sturrock & Stern 1980). The rotational energy can be converted to the thermal energy by Galactic dynamo and magnetic reconnection in  $\sim 10^8 \text{ yr}$  (e.g. Parker 1971; Tanuma et al. 1999).

##### 4.1. Implication for Observations

In our numerical simulations, the magnetic Reynolds number is much smaller than actual value in Galaxy. The Alfvén velocity, resistivity, and magnetic Reynolds number are  $v_A \sim 6 \times 10^6 (B/10 \mu\text{G})(n/0.1 \text{ cm}^{-3})^{-1/2} \text{ cm s}^{-1}$ ,  $\eta \sim 10^7 (T/10^4 \text{ K})^{-3/2} \text{ cm}^2 \text{ s}^{-1}$ , and  $R_m \sim \lambda v_A / \eta \sim 2 \times 10^{20} (B/10 \mu\text{G})(n/0.1 \text{ cm}^{-3})^{-1/2} (\lambda/100 \text{ pc})(T/10^4 \text{ K})^{3/2}$ , if we assume the length of current sheet which is getting thin by the tearing instability, where the local magnetic field strength, number density, and temperature are  $\lambda \sim 100 \text{ pc}$ ,  $B \sim 10 \mu\text{G}$ ,  $n \sim 0.1 \text{ cm}^{-3}$ , and  $T \sim 10^4 \text{ K}$ , respectively. Sweet-Parker current-sheet thickness is

$$\begin{aligned} l_{\text{SP}} &\sim \lambda R_m^{-1/2} \\ &\sim 2 \times 10^{10} \left( \frac{B}{10 \mu\text{G}} \right)^{-1/2} \left( \frac{n}{0.1 \text{ cm}^{-3}} \right)^{1/4} \\ &\quad \left( \frac{\lambda}{100 \text{ pc}} \right)^{1/2} \left( \frac{T}{10^4 \text{ K}} \right)^{-3/4} \text{ cm}. \end{aligned} \quad (34)$$

The time scale of Sweet-Parker reconnection is

$$\begin{aligned}\tau_{\text{SP}} &\sim \frac{\lambda}{v_{\text{in}}} \sim \frac{\lambda}{v_A R_m^{-1/2}} \\ &\sim 3 \times 10^{16} \left(\frac{\lambda}{100 \text{ pc}}\right)^{3/2} \left(\frac{B}{10 \mu\text{G}}\right)^{-1/2} \\ &\quad \left(\frac{n}{0.1 \text{ cm}^{-3}}\right)^{-3/4} \left(\frac{T}{10^4 \text{ K}}\right)^{3/4} \text{ yr} \quad (35)\end{aligned}$$

(under classical Spitzer resistivity), so that the Sweet-Parker type reconnection can not play an important role as energy-release mechanism in actual interstellar medium.

On the other hand, the time scales of the secondary tearing instability and current-sheet thinning are

$$\tau_{2,\text{tear}} \sim \left(\frac{l^3}{\eta v_A}\right)^{1/2} \sim \left(\frac{l}{\lambda}\right)^{3/2} \left(\frac{\lambda}{v_A}\right) R_m^{1/2}, \quad (36)$$

$$\begin{aligned}\tau_{\text{thin}} &\sim \frac{\lambda}{v_A} \\ &\sim 2 \times 10^6 \left(\frac{\lambda}{100 \text{ pc}}\right) \\ &\quad \left(\frac{B}{10 \mu\text{G}}\right)^{-1} \left(\frac{n}{0.1 \text{ cm}^{-3}}\right)^{1/2} \text{ yr}, \quad (37)\end{aligned}$$

respectively. The secondary tearing instability occurs faster than the current-sheet thinning if  $\tau_{2,\text{tear}} < \tau_{\text{thin}}$ , i.e., the current-sheet thickness decreases to

$$\begin{aligned}l &< l_{2,\text{tear}} \\ &= \lambda R_m^{-1/3} \\ &\sim 5 \times 10^{13} \left(\frac{B}{10 \mu\text{G}}\right)^{-1/3} \left(\frac{n}{0.1 \text{ cm}^{-3}}\right)^{1/6} \\ &\quad \left(\frac{\lambda}{100 \text{ pc}}\right)^{2/3} \left(\frac{T}{10^4 \text{ K}}\right)^{-1/2} \text{ cm}. \quad (38)\end{aligned}$$

In this situation, the aspect ratio becomes  $\lambda/(2l_{\text{tear}}) \sim 4 \times 10^6$ . Hence, the secondary tearing instability starts before Sweet-Parker current sheet develops completely. The time scale and wavelength of the secondary tearing instability are calculated to be

$$\begin{aligned}\tau_{2,\text{tear}} &\sim 2 \times 10^6 \left(\frac{B}{10 \mu\text{G}}\right)^{-1/3} \left(\frac{n}{0.1 \text{ cm}^{-3}}\right)^{1/6} \\ &\quad \left(\frac{\lambda}{100 \text{ pc}}\right)^{2/3} \left(\frac{T}{10^4 \text{ K}}\right)^{-1/2} \text{ yr}. \quad (39)\end{aligned}$$

$$\begin{aligned}\lambda_{2,\text{tear}} &\sim 6 \left(\frac{v_A l_{2,\text{tear}}}{\eta}\right)^{1/4} l_{2,\text{tear}} \\ &\sim 6 \lambda R_m^{-1/6} \\ &\sim 1 \times 10^{18} \left(\frac{B}{10 \mu\text{G}}\right)^{-1/6} \left(\frac{n}{0.1 \text{ cm}^{-3}}\right)^{1/12} \\ &\quad \left(\frac{\lambda}{100 \text{ pc}}\right)^{5/6} \left(\frac{T}{10^4 \text{ K}}\right)^{-1/4} \text{ cm}, \quad (40)\end{aligned}$$

respectively.

When a plasmoid moves along the current sheet, the inflow velocity increases behind it so that the current-sheet width decreases to

$$\begin{aligned}l_{\text{min}} &\sim \frac{\eta}{v_{\text{in}}} \\ &\sim \lambda R_m^{-5/6} \\ &\sim 3 \times 10^3 \left(\frac{B}{10 \mu\text{G}}\right)^{-5/6} \left(\frac{n}{0.1 \text{ cm}^{-3}}\right)^{5/12} \\ &\quad \left(\frac{\lambda}{100 \text{ pc}}\right)^{1/6} \left(\frac{T}{10^4 \text{ K}}\right)^{15/12} \text{ cm}, \quad (41)\end{aligned}$$

by assuming  $\lambda_{2,\text{tear}} v_{\text{in}} = l_{2,\text{tear}} v_A$ , where  $v_{\text{in}}$  is the inflow velocity toward the dissipation region. In this transient large inflow stage, the anomalous resistivity can set in, which eventually leads to Petschek type reconnection, since the condition of anomalous resistivity is that the current-sheet thickness must be smaller than the ion Larmor radius [ $\sim 10^7 (T/10^4 \text{ K})^{1/2} (B/10 \mu\text{G})^{-1} \text{ cm}$ ; figure 23]. The current-sheet thickness, otherwise, decreases to the ion Larmor radius through third, forth, and fifth tearing instabilities (figure 23). The third tearing instability occurs when the current-sheet thickness becomes  $\sim 1 \times 10^{12} \text{ cm}$  (wavelength is  $\sim 7 \times 10^{15} \text{ cm}$ ; the time scale of the third tearing instability is  $\sim 10^4 \text{ yr}$ ) in the current sheet generated by the secondary tearing instability. The forth, fifth, sixth, seventh, and eighth tearing instabilities start when the current-sheet thickness becomes  $\sim 4 \times 10^{11} \text{ cm}$ ,  $\sim 2 \times 10^{10} \text{ cm}$ ,  $\sim 1 \times 10^9 \text{ cm}$ ,  $\sim 7 \times 10^7 \text{ cm}$ , and  $\sim 4 \times 10^6 \text{ cm}$ , respectively. Total released magnetic energy is

$$E \sim \left(\frac{B^2}{8\pi}\right) \lambda^3 \sim 10^{50} \left(\frac{B}{10 \mu\text{G}}\right)^2 \left(\frac{\lambda}{100 \text{ pc}}\right)^3 \text{ erg}, \quad (42)$$

if we assume the volume of space filled with the magnetic field is  $\lambda^3$ . Petschek reconnection continues for

$$\begin{aligned}\tau_P &\sim \frac{\lambda}{\epsilon v_A} \\ &\sim 10^7 \left(\frac{\epsilon}{0.1}\right)^{-1} \left(\frac{\lambda}{100 \text{ pc}}\right) \\ &\quad \left(\frac{B}{10 \mu\text{G}}\right)^{-1} \left(\frac{n}{0.1 \text{ cm}^{-3}}\right)^{1/2} \text{ yr}, \quad (43)\end{aligned}$$

which is equal to one-third of the time of Galactic rotation. The magnetic energy-release rate is

$$\begin{aligned}\frac{E}{\tau_P} &\sim 10^{34.5} \left(\frac{\epsilon}{0.1}\right)^{-1} \left(\frac{\lambda}{100 \text{ pc}}\right)^2 \\ &\quad \left(\frac{B}{10 \mu\text{G}}\right)^3 \left(\frac{n}{0.1 \text{ cm}^{-3}}\right)^{-1/2} \text{ erg s}^{-1}. \quad (44)\end{aligned}$$

#### 4.2. Generation of X-ray Gas in Galaxy

Figure 24 shows the 2D distribution of the thermal radiation [ $F \propto n^2 \Lambda(T)$ ], for the typical model (Model A1), where  $\Lambda(T)$  is the cooling function ( $\sim T^a$ ). The above  $a$  is 0.5, -0.6, 2.9, and 1.5, respectively when the unit of temperature is  $T_0 \sim 10^7$  K (figure 24[a]),  $\sim 10^{5-6}$  K (figure 24[b]),  $10^4$  K (figure 24[c]), and  $10^3$  K (figure 24[d]). The X-ray thermal emission shown in figures 24(a) and 24(b) may be observed by *Newton* and *Chandra* in the galaxies.

The bright thermal X-ray emission are observed along the Galactic plane ( $L_X \sim 10^{38}$  erg s $^{-1}$ ; Galactic Ridge X-ray Emission=GRXE; e.g. Kaneda et al. 1997; Koyama et al. 1986a; Makishima 1994; Yamasaki et al. 1997; Yamauchi et al. 1996; Valinia & Marshall 1998). If the magnetic reconnection occurs in the magnetic field of  $B \sim 30$   $\mu$ G, which is generated locally by Galactic dynamo, the hot ( $T \sim 10^8$  K) component ( $n \sim 3 \times 10^{-3}$  cm $^{-3}$ ; Kaneda et al. 1997) of GRXE can be generated (Makishima 1994; Tanuma et al. 1999). After then, the hot gas is confined by the locally strong magnetic field for  $\tau_{\text{cond}} \sim 10^5 (\lambda_{\text{eff}}/1 \text{ kpc}) (T/10^8 \text{ K})^{-1/2}$  yr, where  $\lambda_{\text{eff}}$  is the effective length of helical magnetic field. Note that the radiative cooling time is  $\tau_{\text{rad}} \sim 10^{10.5} (T/10^8 \text{ K}) (n/0.003 \text{ cm}^{-3})^{-1} [\Lambda(10^8 \text{ K})/10^{-23} \text{ erg cm}^3 \text{ s}^{-1}]^{-1}$  yr, which is too much longer than  $\tau_{\text{cond}}$  so that the radiation is negligible, where  $\Lambda(T)$  is the cooling function. The heating time by Petschek type reconnection is  $\tau_P \sim 5 \times 10^5 (\lambda/100 \text{ pc}) (B/30 \mu\text{G})^{-1} (n/0.003 \text{ cm}^{-3})^{1/2}$  yr, so that the heating is balanced with the conduction cooling, where  $\lambda$  is the thickness of the reconnecting magnetic field.

The hot gas are also observed in Galactic halo (Pietz et al. 1998), Galactic center (e.g. Koyama 1989), and clusters of galaxies (e.g. Sarazin 1986). The magnetic reconnection can generate the hot gas (Zimmer, Lesch, & Birk 1997; Birk, Lesch, & Neukirch 1998; Kerp, Lesch, & Mack 1994; Kerp et al. 1996; Soker 1997) and confine it in the magnetic loop for a long time.

The reconnection jet is accelerated in opposite directions by the magnetic tension force, which is like bipolar-jet. Inside the helical tube and current sheet, the temperature and gas pressure increases by the self-pinch of helical magnetic tube or the collision between the reconnection jet and interstellar cloud so that the star formation may be triggered.

#### 5. SUMMARY

We examined the magnetic reconnection triggered by a supernova (or a point explosion) in interstellar medium, by performing two-dimensional resistive magnetohydrodynamic (MHD) numerical simu-

lations with high resolution and large simulation region size. We found that the magnetic reconnection starts long after a supernova shock (fast-mode MHD shock) passes a current sheet. The long current sheet evolves as follows: (i) The supernova shock perturbs the current sheet. The magnetic reconnection does not occur immediately after the passage of the supernova shock across the current sheet, because the magnetic field lines do not have enough time to reconnect. The tearing-mode instability is excited by the supernova shock, and the current sheet becomes thin in its nonlinear stage. (ii) The current-sheet thinning is saturated when the current-sheet thickness becomes comparable to that of Sweet-Parker current sheet. After that, Sweet-Parker type reconnection starts. (iii) The current-sheet length increases during the Sweet-Parker type reconnection, because the gas in the Sweet-Parker current sheet flows out along the current sheet. The secondary tearing mode instability occurs in the Sweet-Parker current sheet, because the ratio of length to thickness of the current sheet exceeds  $\sim 100$ . (iv) Further current-sheet thinning occurs after the islands moved away. The anomalous resistivity sets in, and Petschek type reconnection starts. The magnetic energy is released while the magnetic islands are moving during the Petschek type reconnection. The phenomena depend only on  $\beta$  (i.e.  $B$ ), not on the distance between the supernova and current sheet, nor the energy of initial explosion.

In actual Galaxy, the magnetic reconnection can occur when the non-parallel magnetic fields (i.e. current sheet) exist and a point explosion such as supernova, superbubble, collision of the clouds or stellar wind will occur near the current sheet. The initial current sheet will be made by the turbulence, Parker(1966) instability (Tanuma et al. 1998), magnetorotational instability (Balbus & Hawley 1991), collision of galaxies etc. The magnetic islands (or helical tubes) confine the hot gas, and flow fast along the current sheet. The interstellar gas flows also inside the magnetic helical tube in three-dimensional (3D) view. The gas pressure is very high inside the magnetic islands so that the gas flows along the tube actually. In the magnetic tube and current sheet, the temperature and gas pressure are very high so that the star forming region may appear. We will apply the results to the X-rays from the Galactic ridge, Galactic halo, Galactic center and clusters of galaxies. We will examine the reconnection in 3D simulations in future works.

The authors thank K. Makishima in University of Tokyo, R. Matsumoto in University of Chiba, and T. Magara in Kyoto University for various fruitful discussion. The numerical computations were carried

out on VPP300/16R and VX/4R at the National As-

tronomical Observatory of Japan.

## REFERENCES

- Balbus, S. A. & Hawley, J. F. 1991, *ApJ*, 376, 214  
 Beck, R., Brandenburg, A., Moss, D., Shukurov, A., & Sokoloff, D. 1996, *ARA&A*, 34, 155  
 Birk, G. T., Lesch, H., & Neukirch, T. 1998, *MNRAS*, 296, 165  
 Biskamp, D. 1993, *Nonlinear Magnetohydrodynamics* (New York: Cambridge Univ. Press), ch 6.5  
 Blitz, L. et al. 1999, *ApJ*, 514, 818  
 Fu, Z. F. & Lee, L. C. 1986, *J. Geophys. Res.*, 91, 13373  
 Furth, H. P., Killeen, J., & Rosenbluth, M. N. 1963, *Phys. of Fluid*, 6, 459  
 Habe, A. & Ikeuchi S. 1980, *Prog. Theor. Phys.*, 64, 1995  
 Hanasz, M. & Lesch, H. 1998, *A&A*, 332, 77  
 Hayashi, M. R., Shibata, K., & Matsumoto, R. 1996, *ApJ*, 468, L37  
 Heiles et al. 1976, *ARA&A*, 14, 1  
 Hoshino, M., Mukai, T., Yamamoto, T., & Kokubun, S. 1998, *JGR*, 103, 4509  
 Ikeuchi, S. 1988, *Fund. Cosmic Phys.*, 12, 255  
 Kaneda, H., Makishima, K., Yamauchi, S., Matsuzaki, K., & Yamasaki, N. 1997, *ApJ*, 491, 638  
 Kerp, J., Lesch, H., & Mack, K.-M. 1994, *A&A*, 286, L13  
 Kerp, J. et al. 1996, *A&A*, 312, 67  
 Koyama, K., Makishima, K., Tanaka, Y., & Tsunemi, H., 1986, *PASJ*, 38, 121  
 Koyama, K., Ikeuchi, S., & Tomisaka, K. 1986, *PASJ*, 38, 503  
 Koyama, K. et al. 1989, *Nature*, 339, 603  
 Li, F. & Ikeuchi, S. 1992, *ApJ*, 390, 405  
 Makishima, K. 1994, in *Elementary Processes in Dense Plasmas*, ed. S. Ichimaru & S. Ogata (Reading: Addison and Wesley), 47  
 Magara, T. & Shibata, K. 1999, *ApJ*, 514, 456  
 Norman, C. A. & Ikeuchi, S. 1989, *ApJ*, 345, 372  
 Odstrčil, D. & Karlický, M. 1997, *A&A*, 326, 1252  
 Ohno, H. & Shibata, S. 1993, *MNRAS*, 262, 953  
 Parker, E. N. 1957, *J. Geophys. Res.*, 62, 509  
 Parker, E. N. 1966, *ApJ*, 145, 811  
 Parker, E. N. 1971, *ApJ*, 163, 255  
 Parker, E. N. 1992, *ApJ*, 401, 137  
 Petschek, H. E. 1964, in *AAS-NASA Symp. on the Physics of solar flares*, ed. W. N. Hess (Washington DC: NASA), 425  
 Pietz, J., Kerp, J., Kalberla, P. M. W., Burton, W. B., Hartmann, D., & Mebold, U. 1999, *A&A*, 332, 55  
 Priest, E. R. & Forbes, T. G. 1986, *JGR*, 91, 5579  
 Rand, R. J. & Kulkarni, S. R. 1989, *ApJ*, 343, 760  
 Sarazin, C. L. 1986, *Rev. of Mod. Phys.*, 58, 1  
 Sato, T. & Hayashi, T. 1979, *Phys. Fluids*, 22, 1189  
 Shibata, K. 1996, *Adv. Space Res.*, 17, 9  
 Shibata, K. 1997, in *SOHO 5th Conference, The Corona and Solar Wind Near Minimum Activity*, ed. A. Wilson ( : ESA), 103  
 Shibata, K. & Yokoyama, T. 1999, *ApJ*, 526, L49  
 Sofue, Y., Klein, U., Beck, R., & Wielebinski, R. 1986, *ARA&A*, 24, 456  
 Soker, N. 1997, *ApJ*, 488, 572  
 Spitzer, L. 1962, *Physics of Fully Ionized Gases* (New York: Interscience)  
 Steinolfson, R. S. & van Hoven, G. 1983, *Phys. Fluid*, 261, 117  
 Steinolfson, R. S. & van Hoven, G. 1984, *Phys. Fluid*, 27, 1207  
 Sturrock, P. A. & Stern, R. 1980, *ApJ*, 238, 98  
 Sweet, P. A. 1958, in *IAU Symp. 6, Electromagnetic Phenomena in Cosmological Physics*, ed. B. Lehnert (Cambridge: Cambridge Univ. Press), 123  
 Tajima, T. & Shibata, K. 1997, *Plasma Astrophysics* (Reading: Addison and Wesley), ch 5  
 Tanuma, S. et al. 1998, in *Numerical Astrophysics 1998*, ed. S. M. Miyama, T. Hanawa, & K. Tomisaka (Dordrecht: Kluwer Academic Publishers), 119  
 Tanuma, S., Yokoyama, T., Kudoh, T., Matsumoto, R., Shibata, K., & Makishima, K. 1999, *PASJ*, 51, 161  
 Tenorio-Tangle, G. & Bodenheimer, P. 1988, *ARA&A*, 26, 145  
 Tomisaka, K. 1998, *MNRAS*, 298, 797  
 Tsuneta, S. 1996, *ApJ*, 456, 63  
 Ugai, M. 1986, *Phys. of Fluid*, 29, 3659  
 Ugai, M. 1992, *Phys. of Fluid*, 4, 2953  
 Valinia, A., Tajima, T., Matsumoto, R., Makishima, K., & Shibata, K. 1996, *ASP Conf. Ser. 88, Clusters, Lensing and the Future of the Universe*, ed. V. Trimble & A. Reisenegger, 205  
 Valinia, A. & Marshall, F. E. 1998, *ApJ*, 505, 134  
 Vallée, J. P. 1997, *Fund. Cosmic Phys.*, 19, 1  
 Yamasaki, N. Y et al. 1997, *ApJ*, 481, 821  
 Yamauchi, S., Kaneda, H., Koyama, K., Makishima, K., Matsuzaki, K., Sonobe, T. Tanaka, Y., & Yamasaki, N. 1996, *PASJ*, 48, L15  
 Yokoyama, T. & Shibata, K. 1995, *Nature*, 375, 42  
 Yokoyama, T. & Shibata, K. 1997, *ApJ*, 474, L61  
 Zimmer, F., Lesch, H., & Birk, G. T. 1997, *A&A*, 320, 746

FIG. 1.— The situation of our problem. The interstellar space is filled with the magnetic fields. The magnetic fields are not exactly parallel to one another, and have numerous current sheets at all scales. We examine how the magnetic reconnection occurs when a supernova perturbs the current sheet. We calculate the interaction between the supernova and current sheet by assuming simulation region shown by black square.

FIG. 2.— Initial condition of the typical model (Model A1) of our numerical simulations. We assume the uniform temperature and uniform total pressure ( $P_g + B^2/8\pi$ ) everywhere. The current-sheet thickness is  $2l^{\text{init}}H = 2H$  where  $H$  is the unif of length. The units of velocity and time are sound velocity ( $C_s$ ) and  $H/C_s$ , respectively. We solve the non-dimensional equations numerically in this paper. The magnetic field is assumed as  $B_x = B_0 \tanh(y/H)$  and  $B_y = 0$ . The plasma  $\beta$  is  $8\pi p_0/B_0^2 = 0.2$  everywhere outside the current sheet. The distance between the supernova and current sheet is  $y_{\text{ex}}H = 7H$ . Minimum grid size is  $(\Delta x, \Delta y) \geq (0.20H, 0.025H)$ . The number of grid points is  $(465 \times 602)$ . The simulation region is  $-120.9H \leq x \leq 120.9H$  and  $-54.8H \leq y \leq 81.7H$ . We assume no gravity ( $g = 0$ ).

FIG. 3.— Time evolution of 2D distribution of the current density ( $J$ ), density ( $\rho$ ), and gas pressure ( $p_g$ ), with the magnetic field lines, in an early phase for the typical model (Model A1). The half-thickness of initial current sheet is  $l^{\text{init}} = 1$  ( $|y| < 1$ ). The initial density, gas pressure, and magnetic field outside the current sheet are  $n_0 = 1$ ,  $p_0 = 0.6$ , and  $B_0 \sim 8.7$ , respectively. The plasma  $\beta$  is  $8\pi p_0/B_0^2 = 0.2$  ( $|y| > 1$ ). The Alfvén velocity is  $B_0/(4\pi\rho)^{-1/2} \sim 2.5$ . We put a supernova at  $(0, y_{\text{ex}}) = (0, 7)$ . The supernova expands in the magnetic field. These figures show the expanding supernova shock well. The slow shock propagates along the magnetic field. The shock front is in a high-density region, while the center of supernova is in a low-density region.

FIG. 4.— Time evolution of 2D distribution of the temperature ( $T$ ), magnetic field lines, and velocity vectors, in a main phase of the magnetic reconnection triggered by a supernova, for the same model shown in figure 2 (Model A1). A supernova appears at  $(x, y) = (0, 7)$ , and disturbs the current sheet. The tearing instability occurs, and the current sheet gets thin ( $t \sim 100 - 200$ ). Sweet-Parker type reconnection occurs at  $t \sim 200 - 250$ . Petschek type reconnection starts at  $t \sim 270$ . The magnetic reconnection heats the gas, and accelerates it by the magnetic tension force. The hot gas of supernova flows into the diffusion region. The magnetic islands are generated in the current sheet sequentially. The heated gas is confined in the magnetic island.

FIG. 5.— Time evolution of 2D distribution of the gas pressure ( $p_g$ ), magnetic field lines, and velocity vectors, in a main phase of the magnetic reconnection triggered by a supernova, for the same model shown in figure 2 (Model A1). The initial gas pressure is  $p_g = p_0 + (p_0/\beta) \cosh^{-2}(y)$  where  $p_0 = 0.6$  and  $\beta = 0.2$ . The gas pressure is high in the current sheet, especially inside magnetic islands ( $t \sim 270, 300$ ).

FIG. 6.— Time evolution of 2D distribution of the current density ( $J$ ), magnetic field lines, and velocity vectors, in a main phase of the magnetic reconnection triggered by a supernova, for the same model shown in figure 2 (Model A1). The initial magnetic field is  $B_x = B_0 \tanh(y)$  and  $B_y = 0$  where  $B_0 = 8.58$ . The plasma  $\beta$  is  $p_0/(B_0^2/8\pi) = 0.2$ . The initial current-sheet thickness is  $2l^{\text{init}} = 2$ . The initial current density is  $\sim B_0/2 \sim 4.5$  in the current sheet. The current density increases in the diffusion region. Petschek type reconnection generates two standing slow shocks in the current sheet (see also figure 15). The slow shocks appear also behind the moving magnetic islands (see the figures at  $t = 270$  and  $300$ ).

FIG. 7.— The schematic illustration of the results of our simulation (Model A1). A supernova disturbs the current sheet (figure a). The tearing instability occurs long after the passage of fast shock wave (figure b). The current-sheet thickness decreases. Sweet-Parker type reconnection occurs (figure c). The secondary tearing instability occurs in the Sweet-Parker current sheet (figure d). A magnetic island is generated (figure e). Petschek type reconnection starts after the islands moves away (figure f). It heats the interstellar gas, and accelerates the heated gas by the magnetic tension force to the Alfvén velocity. Many islands are generated sequentially, and confine the heated gas.

FIG. 8.— Time variation of the half-thickness of the current sheet ( $l$ ), which is defined by the minimum half-width of half-maximum of the gas pressure calculated parallel with  $y$ -axis in  $|x| < 25$ , for the same model shown in figure 2 (Model A1). The supernova shock wave perturbs the current sheet ( $t < 2$ ). The magnetic field dissipates slowly ( $t < 100$ ). The tearing instability occurs ( $t \sim 100 - 200$ ). The current-sheet thinning occurs during the tearing instability. Sweet-Parker type reconnection occurs ( $t \sim 200 - 250$ ). The secondary tearing instability occurs in the Sweet-Parker current sheet ( $t \sim 250 - 270$ ). The current-sheet thickness decreases again by the secondary tearing instability. Petschek type reconnection occurs ( $t \sim 270 - 400$ ). The current sheet gets much thin during the Petschek type reconnection. At  $t \sim 390 - 400$ , the current sheet is very thin because the magnetic island is moving.

FIG. 9.— The schematic illustration of the time variation of the current-sheet thickness. At first, the magnetic field dissipates slowly. Second, the current-sheet thinning occurs by the tearing instability in a nonlinear phase. Sweet-Parker type reconnection occurs in the thin current sheet. Next, the secondary tearing instability occurs in the Sweet-Parker current sheet. The current-sheet thickness decreases more. Petschek type reconnection starts. The solid line moves to upper side for large- $\beta$ , small- $y_{\text{ex}}$ , and large- $p_{\text{ex}}$  models, while it moves to lower side for small- $\beta$ , large- $y_{\text{ex}}$ , and small- $p_{\text{ex}}$  models.

FIG. 10.— Time variation of the magnetic flux, for the same model shown in figure 2 (Model A1). The magnetic flux is defined by  $\psi(t) = \int_0^t \max_{|x| < 10} |(B_x v_y)[\text{at}(x, 2)] + (B_x v_y)[\text{at}(x, -3)]| dt$ . The magnetic flux increases during the tearing instability. The magnetic flux grows by  $\log \psi(t) \sim 0.00257t$ , which is shown by the dashed line. The rate is 1/2 times the growth rate of the linear phase (Magara & Shibata 1999).

FIG. 11.— Time variation of the inflow velocity toward the diffusion region, for the same model shown in figure 2 (Model A1). Figure (a) shows the inflow velocity, defined by  $v_{\text{in}} = \max_{|x| < 10} [v_y[\text{at}(x, -3)] - v_y[\text{at}(x, 2)]] / 2$ . Figure (b) shows the Alfvén velocities at  $(x_0, 2)$  (dotted line) and  $(x_0, -3)$  (solid line), where  $x_0$  is the point where maximum inflow velocity is attained. Figure (c) shows the reconnection rate defined by  $\epsilon = v_{\text{in}} / v_{\text{A},s}$ , where  $v_{\text{A},s}$  is smaller one between  $v_{\text{A}}[\text{at}(x_0, -2)]$  and  $v_{\text{A}}[\text{at}(x_0, 2)]$ . The inflow velocity and reconnection rate are  $v_{\text{in}} \sim 0.1 - 0.2$  and  $\epsilon \sim 0.05 - 0.1$  during Petschek type reconnection, while they are  $\sim 0.05$  and  $\sim 0.02$  during Sweet-Parker type reconnection. At the beginning of Sweet-Parker type reconnection ( $t \sim 200 - 220$ ), they increase to  $\sim 0.16$  and  $\sim 0.6$ , respectively, because of the nonlinear growth of the tearing instability (see the arrow).

FIG. 12.— Time variation of the various energies, showing how the energy conversion occurs by the magnetic reconnection in a central region ( $|x| < 25$  and  $|y| < 3$ ) for the same model shown in figure 2 (Model A1). The magnetic energy is converted to the thermal energy slowly by Sweet-Parker type reconnection and secondary tearing instability ( $t \sim 200 - 270$ ), and quickly by Petschek magnetic reconnection ( $t \sim 270 - 360$ ). The magnetic islands suppress the magnetic reconnection when they are staying at the reconnection region ( $t \sim 310$  and  $360$ ). The magnetic energy-release rate increases to  $\sim 40$  at the beginning of Sweet-Parker reconnection because the inflow velocity increases (see the arrow). Maximum energy release rate is  $-dE_{\text{mag}}/dt \sim 70 - 80$  during Petschek type reconnection ( $t \sim 340$ ). The non-dimensional thermal energy of the initial supernova is  $E_{\text{ex}} \sim 1800$  in the numerical simulation. Main thermal energy is supplied from the magnetic energy, not from the initial supernova.

FIG. 13.— The schematic illustration of the time variation of the magnetic energy-release rate in the reconnection region. The magnetic dissipation occurs slowly at first. The tearing instability starts much after the passage of supernova shock. Sweet-Parker type reconnection occurs after current-sheet thinning by the nonlinear phase of the tearing instability. The magnetic energy-release rate increases once at  $t \sim 200 - 220$  (see the arrow) because the inflow velocity increases. After then, the magnetic energy is released slowly by Sweet-Parker reconnection. The secondary tearing instability occurs in Sweet-Parker current sheet. The magnetic energy-release is suppressed by the small islands generated by the secondary tearing instability. Petschek type reconnection starts at last. The magnetic energy is converted to the thermal energy violently during Petschek type reconnection.

FIG. 14.— Time variations of (a) the maximum drift velocity ( $v_d = J/\rho$ ), (b) density  $\rho$  where the maximum  $v_d$  is attained, (c) current density ( $J$ ) where the maximum  $v_d$  is attained, and (d) maximum reconnection rate defined by  $\eta J$  ( $-2.6 < y < 1, |x| < 10$ ), for the same model shown in figure 2 (Model A1). The current density increases to  $J \sim B/l_{\text{SP}} \sim 20$  after Sweet-Parker reconnection starts. The density in the current sheet decreases during Sweet-Parker type reconnection and secondary tearing instability ( $t \sim 200 - 270$ ). The drift velocity exceeds the critical value ( $v_c = 100$ ) so that the anomalous resistivity sets in ( $t \sim 270$ ). Petschek type reconnection starts so that the reconnection rate increases.

FIG. 15.— Profile of the variables in  $x = -12$  at  $t \sim 342$ , for the same model shown in figure 2 (Model A1).  $p_g$ ,  $T$ ,  $\rho$ , and  $|v_y|$  are higher in the current sheet than those outside the sheet. The magnetic field strength is lower in the current sheet. The profile of the current density has two peaks ( $y \sim -2.3$  and  $0.7$ ), corresponding to the slow shocks. The density ratio across the shock front is  $X = n_2/n_1 \sim 0.3$ , where subscripts 1 and 2 mean that the values are attained at  $y \sim -2.1$  and  $-0.3$ , respectively. The gas pressure ratio is  $p_{g2}/p_{g1} \sim 0.1$ , while 0.47 is required by Rankine-Hugoniot relation.

FIG. 16.— 2D distribution of the gas pressure at  $t = 160$  (a), 300 (b), and 450 (c), for the same model shown in figure 2 (Model A1). Figures (a), (b), and (c) show the distributions during the tearing instability, a little after Petschek type reconnection starts, and after Petschek type reconnection stops. The gas is ejected from the diffusion region toward the left and right boundaries. The gas pressure increases in the current sheet near the boundaries. It extends the current sheet near the boundaries. Furthermore, the magnetic tension suppresses the magnetic reconnection because the magnetic field is bent outside the current sheet.

FIG. 17.— Time variation of the magnetic energy-release rate ( $|x| < 25, |y| < 3$ ), for Model B1 ( $\beta = 0.1$ ; dotted line), Model A2 ( $\beta = 0.2$ ; solid line), and Model B2 ( $\beta = 0.3$ ; dashed line). The magnetic energy-release rate decreases with  $\beta$ . The maximum magnetic energy release rates are  $-dE_{\text{mag}}/dt \sim 250, 70$ , and  $20$ , respectively.

FIG. 18.— Dependence on  $\beta$  of (a) the maximum energy release rate ( $dE_{\text{mag}}/dt$ ) in the reconnection region ( $|x| < 25, |y| < 3$ ), (b) the time when maximum  $dE_{\text{mag}}/dt$  is attained, (c) maximum temperature of heated gas, and (d) maximum velocity of the reconnection jet (Model A2, B1-5, and E). (a) The magnetic energy-release rate decreases with  $\beta$ . The solid line and dotted line show the maximum magnetic energy release rates by Petschek and Sweet-parker type reconnections, respectively. The rates are determined by Poynting flux entering into the reconnection region. The theoretical  $\beta$ -dependence is shown by the dashed line [ $2(B_0^2/4\pi)S\epsilon v_A^{\text{init}} \propto \beta^{-3/2}$  where we assume  $\epsilon \sim 0.1$  and  $S \sim 26$ ] for Petschek type reconnection, and the dashed-dotted line [ $2(B_0^2/4\pi)SR_{\text{m,SP}}^{-1/2}v_A^{\text{init}} \propto \beta^{-5/4}$  where we assume  $R_{\text{m,SP}} \sim 10^4$  and  $S \sim 67$ ] for Sweet-Parker type reconnection. The results of the uniform-resistivity model (Model E) are also shown by the triangle ( $\triangle$ ). (b) The time scale increases with  $\beta$ . The magnetic phenomena occur in the Alfvén time ( $\tau_A \propto \beta^{1/2}$ ; dashed line) or the time scale of tearing instability ( $\tau_A \propto \beta^{1/4}$ ; dashed-dotted line). (c) The maximum temperature decreases with  $\beta$ . It is determined by  $T_{\text{max}} \propto (1 + 1/\beta)$ . (d) The maximum velocity decreases with  $\beta$ . The gas is accelerated to Alfvén velocity  $v_A \propto \tau_A^{-1} \propto \beta^{-1/2}$ .

FIG. 19.— Dependence of the results on the distance ( $y_{\text{ex}}$ ) between the supernova and current sheet (Models A2 and C1-3; see the caption of figure 18). Figure (a) shows the maximum magnetic energy release rate in the reconnection region ( $|x| < 25$ ,  $|y| < 3$ ). It decreases with  $y_{\text{ex}}$ . Figure (b) shows the time when maximum magnetic energy release rate is attained. It increases with  $y_{\text{ex}}$  because the initial perturbation decreases. Figures (c) and (d) show the maximum temperature of the heated gas and maximum velocity of the reconnection jet. They do not depend on  $y_{\text{ex}}$ .

FIG. 20.— Dependence of the results on the energy of the initial explosion ( $E_{\text{ex}} \propto p_{\text{ex}} \sim p_{\text{ratio}}$ ) (Models A2 and D1-4; see the caption of Figure 18). Figure (a) shows the maximum magnetic energy release rate in the reconnection region ( $|x| < 25$ ,  $|y| < 3$ ). It does not depend on  $p_{\text{ratio}}$ . Figure (b) shows the time when maximum magnetic energy release rate is attained. It does not depend on  $p_{\text{ratio}}$ . Figure (c) shows the maximum temperature of the gas heated by the magnetic reconnection. It increases with  $p_{\text{ratio}}$  because the hot gas flows into the reconnection region. Figure (d) shows the velocity of the reconnection jet. It does not depend on  $p_{\text{ratio}}$  because the velocity of the reconnection jet is determined by Alfvén velocity. The results of two-supernovae[at (0, 7) and (0, -7)]-model (Model D5) are indicated by  $\times$ . The magnetic reconnection starts early in the model because the current sheet is compressed from both sides.

FIG. 21.— Time variations of the half-thickness of the current sheet ( $l$ ), defined by the minimum half-width of half-maximum of the gas pressure in  $|x| < 25$ , for Model A2 ( $\eta_0 = 0.015$ ; thick solid line), Model F1 ( $\eta_0 = 0.03$ ; thin dotted line), Model F2 ( $\eta_0 = 0.0075$ ; thin dashed-dotted line), Model F3 ( $\eta_0 = 0.00375$ ; thin dashed and double-dotted line), Model F4 ( $\eta_0 = 0.001$ ; thin dashed line), Model F5 ( $\eta_0 = 0.0001$ ; thin solid line), and Model F6 ( $\eta_0 = 0$ ; thick dashed line). In Model A2 ( $\eta_0 = 0.015$ ), the tearing instability ( $t \sim 60 - 150$ ) is followed by Sweet-Parker type reconnection ( $t \sim 150 - 200$ ), secondary tearing instability ( $t \sim 200 - 240$ ), and Petschek type reconnection ( $t > 240$ ) (figure 8). In zero-resistivity model (F6), the current sheet does not dissipates. It starts to become thin at  $t \sim 220$ , and Petschek type reconnection occurs by only numerical resistivity (resistivity-like effect by the numerical noise). In Model F1 ( $\eta_0 = 0.03$ ), the current sheet dissipates early ( $t < 100$ ).

FIG. 22.— Time variations of the magnetic energy-release rate in the reconnection region ( $|x| < 25$ ,  $|y| < 3$ ), for Model A1 ( $\Delta x = 0.20$ ; solid line) and Model A2 ( $\Delta x = 0.25$ ; dotted line). The magnetic energy-release rate does not depend on the grid size because the “background uniform resistivity” ( $\eta_0$ ) is enough large to suppress the numerical resistivity. The time scale of phenomena changes because the grid number of the initial supernova changes, and because the perturbation on the current sheet changes.

FIG. 23.— The schematic illustration of the time variation of the current-sheet thickness in actual interstellar medium. Here we assume  $B \sim 10 \mu\text{G}$ ,  $n \sim 0.1 \text{ cm}^{-3}$ ,  $T \sim 10^4 \text{ K}$ , and Spitzer conductivity. The current-sheet thickness decreases to the ion Larmor radius ( $\sim 10^7 \text{ cm}$ ) by the nonlinear phase of secondary tearing instability or by successive tearing instabilities. When it decreases to the radius, Petschek type reconnection can occur in the interstellar medium.

FIG. 24.— 2D distribution of the thermal radiation ( $F \propto n^2 T^a$ ) for the same model shown in figure 2 (Model A1).  $a$  is 0.5, -0.6, 2.9, or 1.5, respectively, when the unit of temperature is assumed to be  $T_0 \sim 10^7 \text{ K}$  (figure [a]),  $\sim 10^{5-6} \text{ K}$  (figure [b]),  $10^4 \text{ K}$  (figure [c]), or  $10^3 \text{ K}$  (figure [d]), The high-temperature gas in the island and current sheet is bright.



TABLE 1

Parameters. We studied the effect of the grid size (Models A1-2), plasma  $\beta$  (Models B1-5), distance  $y_{\text{ex}}$  between the current sheet and supernova (Models C1-3), energy of initial supernova (Models D1-5), resistivity model (Models E and F1-4), and simulation region size (Models G1-2). Typical model is Model A1.

| Model | $\beta^{\text{a}}$ | $y_{\text{ex}}^{\text{b}}$ | $p_{\text{ratio}}^{\text{c}}$ | $\eta_0^{\text{d}}$ | $\alpha^{\text{e}}$ | SN <sup>f</sup> | $\Delta x^{\text{g}}$ | $\Delta y^{\text{h}}$ | $L_x^{\text{i}}$ | $L_y^{\text{j}}$ | $N_x^{\text{k}}$ | $N_y^{\text{l}}$ |
|-------|--------------------|----------------------------|-------------------------------|---------------------|---------------------|-----------------|-----------------------|-----------------------|------------------|------------------|------------------|------------------|
| A1    | 0.20               | 7.0                        | 500                           | 0.01500             | 0.1                 | 1               | 0.20                  | 0.025                 | 241.8            | 136.5            | 465              | 602              |
| A2    | 0.20               | 7.0                        | 500                           | 0.01500             | 0.1                 | 1               | 0.25                  | 0.025                 | 153.2            | 136.5            | 301              | 602              |
| B1    | 0.10               | 7.0                        | 500                           | 0.01500             | 0.1                 | 1               | 0.25                  | 0.025                 | 153.2            | 136.5            | 301              | 602              |
| B2    | 0.15               | 7.0                        | 500                           | 0.01500             | 0.1                 | 1               | 0.25                  | 0.025                 | 153.2            | 136.5            | 301              | 602              |
| B3    | 0.30               | 7.0                        | 500                           | 0.01500             | 0.1                 | 1               | 0.25                  | 0.025                 | 153.2            | 136.5            | 301              | 602              |
| B4    | 0.50               | 7.0                        | 500                           | 0.01500             | 0.1                 | 1               | 0.25                  | 0.025                 | 153.2            | 136.5            | 301              | 602              |
| B5    | 1.00               | 7.0                        | 500                           | 0.01500             | 0.1                 | 1               | 0.25                  | 0.025                 | 153.2            | 136.5            | 301              | 602              |
| C1    | 0.20               | 10.0                       | 500                           | 0.01500             | 0.1                 | 1               | 0.25                  | 0.025                 | 153.2            | 136.5            | 301              | 602              |
| C2    | 0.20               | 14.0                       | 500                           | 0.01500             | 0.1                 | 1               | 0.25                  | 0.025                 | 153.2            | 136.5            | 301              | 602              |
| C3    | 0.20               | 28.0                       | 500                           | 0.01500             | 0.1                 | 1               | 0.25                  | 0.025                 | 153.2            | 136.5            | 301              | 602              |
| D1    | 0.20               | 7.0                        | 50                            | 0.01500             | 0.1                 | 1               | 0.25                  | 0.025                 | 153.2            | 136.5            | 301              | 602              |
| D2    | 0.20               | 7.0                        | 100                           | 0.01500             | 0.1                 | 1               | 0.25                  | 0.025                 | 153.2            | 136.5            | 301              | 602              |
| D3    | 0.20               | 7.0                        | 250                           | 0.01500             | 0.1                 | 1               | 0.25                  | 0.025                 | 153.2            | 136.5            | 301              | 602              |
| D4    | 0.20               | 7.0                        | 1000                          | 0.01500             | 0.1                 | 1               | 0.25                  | 0.025                 | 153.2            | 136.5            | 301              | 602              |
| D5    | 0.20               | $\pm 7.0$                  | 500                           | 0.01500             | 0.1                 | 2               | 0.25                  | 0.025                 | 153.2            | 136.5            | 301              | 602              |
| E     | 0.20               | 7.0                        | 500                           | 0.01500             | 0.0                 | 1               | 0.25                  | 0.025                 | 153.2            | 136.5            | 301              | 602              |
| F1    | 0.20               | 7.0                        | 500                           | 0.03000             | 0.0                 | 1               | 0.25                  | 0.025                 | 153.2            | 136.5            | 301              | 602              |
| F2    | 0.20               | 7.0                        | 500                           | 0.00750             | 0.1                 | 1               | 0.25                  | 0.025                 | 153.2            | 136.5            | 301              | 602              |
| F3    | 0.20               | 7.0                        | 500                           | 0.00375             | 0.1                 | 1               | 0.25                  | 0.025                 | 153.2            | 136.5            | 301              | 602              |
| F4    | 0.20               | 7.0                        | 500                           | 0.00100             | 0.1                 | 1               | 0.25                  | 0.025                 | 153.2            | 136.5            | 301              | 602              |
| F5    | 0.20               | 7.0                        | 500                           | 0.00010             | 0.1                 | 1               | 0.25                  | 0.025                 | 153.2            | 136.5            | 301              | 602              |
| F6    | 0.20               | 7.0                        | 500                           | 0.00000             | 0.0                 | 1               | 0.25                  | 0.025                 | 153.2            | 136.5            | 301              | 602              |
| G1    | 0.50               | 7.0                        | 500                           | 0.01500             | 0.1                 | 1               | 0.25                  | 0.025                 | 153.2            | 66.7             | 301              | 349              |
| G2    | 0.50               | 7.0                        | 500                           | 0.01500             | 0.1                 | 1               | 0.25                  | 0.025                 | 153.2            | 44.9             | 301              | 270              |

<sup>a</sup>the ratio of the gas to magnetic pressure outside the current sheet

<sup>b</sup>the distance between the supernova and current sheet

<sup>c</sup>the initial gas pressure of the supernova

<sup>d</sup>“background uniform resistivity”

<sup>e</sup>the parameter of anomalous resistivity; 0.1 for the anomalous resistivity, and 0.0 for the uniform resistivity

<sup>f</sup>number of the initial supernovae

<sup>g</sup>minimum grid size in  $x$ -axis

<sup>h</sup>minimum grid size in  $y$ -axis

<sup>i</sup>simulation region size in  $x$ -axis

<sup>j</sup>simulation region size in  $y$ -axis

<sup>k</sup>number of grid points in  $x$ -axis

<sup>l</sup>number of grid points in  $y$ -axis

This figure "f01.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0009088v2>

This figure "f02.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0009088v2>

This figure "f03.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0009088v2>

This figure "f04.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0009088v2>

This figure "f05.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0009088v2>

This figure "f06.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0009088v2>

This figure "f07.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0009088v2>



This figure "f08.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0009088v2>

This figure "f09.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0009088v2>

This figure "f10.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0009088v2>

This figure "f11.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0009088v2>

This figure "f12.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0009088v2>

This figure "f13.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0009088v2>

This figure "f14.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0009088v2>

This figure "f15.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0009088v2>



This figure "f16.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0009088v2>

This figure "f17.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0009088v2>

This figure "f18.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0009088v2>

This figure "f19.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0009088v2>

This figure "f20.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0009088v2>

This figure "f21.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0009088v2>

This figure "f22.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0009088v2>

This figure "f23.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0009088v2>



This figure "f24.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0009088v2>